arXiv:1110.4082v2 [math.CV] 5 Dec 2011

BOUNDING THE RANK OF HERMITIAN FORMS AND RIGIDITY FOR CR MAPPINGS OF HYPERQUADRICS

DUSTY GRUNDMEIER, JIŘÍ LEBL, AND LIZ VIVAS

Abstract. Using Green's hyperplane restriction theorem, we prove that the rank of a Hermitian form on the space of holomorphic polynomials can be bounded by a constant depending only on the maximum rank of the form restricted to affine manifolds. As an application we prove a rigidity theorem for CR mappings between hyperquadrics in the spirit of the results of Baouendi-Huang and Baouendi-Ebenfelt-Huang. If we have a real-analytic CR mapping of a hyperquadric not equivalent to a sphere to another hyperquadric $Q(A,B)$, then either the image of the mapping is contained in a complex affine subspace or A is bounded by a constant depending only on B. Finally, we prove a stability result about existence of nontrivial CR mappings of hyperquadrics. That is, as long as both A and B are sufficiently large and comparable, then there exist CR mappings whose image is not contained in a hyperplane. The rigidity result also extends when mapping to hyperquadrics in infinite dimensional Hilbert-space.

1. Introduction

A real-valued polynomial, or a real-analytic function, $r(z,\bar{z})$ on $\mathbb{C}^n$ can be regarded as a Hermitian form by considering the matrix of coefficients of the series. That is, in multi-index notation, write

$$r(z,\bar{z}) = \sum_{\alpha\beta} c_{\alpha\beta} z^\alpha \bar{z}^\beta. \tag{1}$$

The matrix $C = [c_{\alpha\beta}]_{\alpha\beta}$ is Hermitian symmetric if and only if r is real-valued (C is uniquely determined by r).

When we apply linear algebra terminology (such as *rank*, *eigenvalues*, *signature*, or *positive semidefinite*) to r, we simply refer to the underlying matrix C. When r is a polynomial, the terminology is obvious. When r is real-analytic, then C is an infinite matrix. After possibly rescaling so that the series converges in a neighborhood of the closed unit polydisc, C defines a Hermitian trace-class operator, and the terminology easily extends to the real-analytic case; see section 5. While the matrix depends on the point where we expand the series, we show that the rank does not change under a biholomorphic change of coordinates. In particular the rank does not depend on the point where we expand the series.

In order to state the first result, we need to define what we mean by the rank of the restriction to an affine manifold. The Grassmannian $G(m,n)$ is the manifold of all m-dimensional linear subspaces of $\mathbb{C}^n$, and the affine Grassmannian $G_{m,n}$ is the manifold of all affine m-dimensional subspaces in $\mathbb{C}^n$. Notice that $G_{m,n}$ is an open set in $G(m+1,n+1)$.

Let $U \subset \mathbb{C}^n$ be a convex neighborhood of the origin, and define $U^* := \{z : \bar{z} \in U\}$. Suppose that the series (1) converges in $U \times U^*$. Let L be an affine m-plane in $\mathbb{C}^n$ (i.e. an

---

Date: December 5, 2011.

The first author acknowledges support from NSF grant DMS 0838434.

The second author was in part supported by NSF grant DMS 0900885.

element of $G_{m,n}$). Let $E\colon \mathbb{C}^m \to \mathbb{C}^n$ be an affine embedding of L in $\mathbb{C}^n$. If L intersects U, then define

$$\operatorname{rank} r|_L := \operatorname{rank} r \circ E. \tag{2}$$

It is not hard to show that the rank does not depend on the particular embedding E, and therefore the notation makes sense. Furthermore as U is convex $U \cap L$ is always connected. If L does not intersect U, then define

$$\operatorname{rank} r|_L := -\infty. \tag{3}$$

Let $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

Theorem 1.1. *Let $n \geq 2$ and let $1 \leq m \leq n-1$. Let $r(z,\bar{z})$ be a real-analytic function defined in a convex neighborhood $U \subset \mathbb{C}^n$ of the origin such that the complexified series converges in $U \times U^*$. If*

$$\sup_{L \in G_{m,n}} \operatorname{rank} r|_L < \infty, \tag{4}$$

then $\operatorname{rank} r < \infty$.

Moreover, there exists a function $R_{m,n}\colon \mathbb{N}_0 \to \mathbb{N}_0$ such that for any such real-analytic r,

$$\operatorname{rank} r \leq R_{m,n}\left(\max_{L \in G_{m,n}} \operatorname{rank} r|_L\right). \tag{5}$$

It is worthwhile to state the theorem for bihomogeneous polynomials. A polynomial $r(z,\bar{z})$ is said to be *bihomogeneous* of *bidegree* (d,d) if $r(tz,\bar{z}) = r(z,t\bar{z}) = t^d r(z,\bar{z})$. When r is a polynomial then the coefficient matrix is simply a finite matrix. The function $R_{m,n}$ in the following theorem is the same as above.

Theorem 1.2. *Let $n \geq 2$, $z \in \mathbb{C}^{n+1}$, and $1 \leq m \leq n-1$. Then there exists a function $R_{m,n}\colon \mathbb{N}_0 \to \mathbb{N}_0$ such that for any bihomogeneous polynomial $r(z,\bar{z})$,*

$$\operatorname{rank} r \leq R_{m,n}\left(\max_{L \in G(m+1,n+1)} \operatorname{rank} r|_L\right). \tag{6}$$

As an application of Theorem 1.1, we prove a rigidity result for mappings of hyperquadrics. The hyperquadric $Q(a,b) \subset \mathbb{C}^{a+b}$ is the set defined by

$$Q(a,b) := \Bigl\{ z : \sum_{j=1}^{a} \lvert z_j \rvert^2 - \sum_{j=a+1}^{a+b} \lvert z_j \rvert^2 = 1 \Bigr\}. \tag{7}$$

When $b = 0$, then $Q(a,0)$ is simply the sphere. Hyperquadrics and spheres are the CR analogues of flat euclidean space from Riemannian geometry. In CR geometry, however, there is no analogue of the Nash embedding theorem. Therefore, a natural question in CR geometry is to study the CR mappings $f\colon U \to Q(A,B)$ for a CR manifold U. In this paper we take an open subset $U \subset Q(a,b)$. We study hyperquadrics $Q(a,b)$ that are not equivalent to the sphere. By a theorem of Lewy [16] a CR function on $Q(a,b)$ extends to a holomorphic function of both sides of $Q(a,b)$. It is therefore enough to consider real-analytic CR mappings or in other words, restrictions of holomorphic mappings. Therefore we study holomorphic mappings of a neighborhood of U in $\mathbb{C}^{a+b}$ to $\mathbb{C}^{A+B}$ that take $U \subset Q(a,b)$ to $Q(A,B)$. See the books [2,5] for more background information.

When considering mappings between hyperquadrics, we need to consider problems of positivity rather than just rank. See [8,9] for more on positivity conditions.

After a linear fractional change of coordinates we can assume that $a > b$ and $A > B$. It is possible that the change of coordinates has a pole on U, but as $Q(a,b)$ is a real hypersurface, there still has to exist a perhaps smaller dense open set $V \subset U$ such that $f \colon V \to Q(A,B)$ is a CR mapping.

The study of CR mappings between spheres has a long history. When the target is also a sphere ($B = 0$) and the codimension $A - a$ is small, then certain strong rigidity results can be proved; see for example [13] and the references within. Forstnerič proved that sufficiently smooth CR mappings of spheres must be rational of degree bounded by a constant depending only on the dimensions involved [10].

When the target dimension is large, there is less rigidity. When both the source and target are spheres, increasing the dimension of the target always adds new CR mappings as long as the dimension is large enough; see [6]. Furthermore, if the source is a sphere and the target is a hyperquadric not equivalent to a sphere, then for large target dimension not only do we always get new mappings, but we get new rational mappings of arbitrarily large degree; see [7].

We therefore concentrate on the case when $b, B \geq 1$. Baouendi and Huang [4] proved that if $b = B$, then f must be a linear embedding. Later, Baouendi, Ebenfelt, and Huang [3] proved that if $B < 2b - 1$, then after a change of coordinates f can be written as

$$(z_1, \dots, z_n) \mapsto (z_1, \dots, z_a, \psi(z), 0, z_{a+1}, \dots, z_{a+b}, \psi(z), 0), \tag{8}$$

where an arbitrary CR mapping ψ and 0 are vector valued functions with the right number of components. In particular, unless $a = A$ and $b = B$, $f(U)$ is contained in a complex hyperplane. The mapping (8) can be written, after an affine change of coordinates, as the identity mapping direct sum an arbitrary CR mapping going into a lower dimensional ambient space.

For any mapping f of hyperquadrics, if $f(U)$ lies in a complex hyperplane, then we apply an affine change of coordinates and obtain a mapping $\tilde{f}$ from U to $Q(A', B') \times \mathbb{C}^k$ for some $A' \leq A$ and $B' \leq B$. We can study the first $A' + B'$ components of $\tilde{f}$ as a mapping from $Q(a,b)$ to $Q(A', B')$, as the last k components of $\tilde{f}$ are arbitrary. Therefore, it is natural to study those mappings where $f(U)$ is not contained in a hyperplane.

Theorem 1.3. *Let $a > b \geq 1$, $U \subset Q(a,b)$ be a connected open set, and $f \colon U \to Q(A,B)$ be a real-analytic CR mapping such that $f(U)$ does not lie in a complex hyperplane, then*

$$A \leq N(a,b,B), \tag{9}$$

where $N = N(a,b,B)$ is a constant depending only on a, b and B.

Note that the hypotheses on $Q(a,b)$ mean that $Q(a,b)$ is not equivalent to a sphere. If $Q(a,b)$ were equivalent to a sphere, there is no rigidity. In this sense, the theorem is optimal.

We are interested in what happens when $B \geq 2b - 1$. In this situation, there do exist nontrivial mappings that do not map to a hyperplane. See sections 8 and 9 for a general method of constructing mappings.

The bound that we obtain on A is not sharp. For example when $a = 2$ and $b = 1$, it is possible to use the method in this paper to obtain an explicit bound

$$A \leq K_{1,3}(B+1) = K_2(K_3(B+1)). \tag{10}$$

See section 3 for the definition of $K_n(k)$. When $B = 1$ we obtain $A \leq 3$, though we know that $A = 2$ by the result of Baouendi-Huang. Therefore, we dispense with finding explicit formulas for $N(a, b, B)$.

Unlike in the sphere to sphere case, there do exist non-rational mappings, and rational mappings of arbitrarily high degree. In Section 9 we prove a stability result about existence of hyperquadric mappings that do not map into hyperplanes as long as both A and B are sufficiently large and comparable. The proof is constructive and all constructed mappings are monomial; that is, all components are single monomials.

The stability result discussed above shows that Theorem 1.3 is optimal in the sense that super-rigidity as studied by Baouendi-Huang and Baouendi-Ebenfelt-Huang only appears in small codimension. In general the best constant N must grow asymptotically at least as a linear function of B. The methods explored in this paper do not readily give the best bound for N as we have mentioned above, but we do obtain that N asymptotically grows no faster than a polynomial in B.

Finally, we also note that the rigidity result extends when the target is an infinite dimensional hyperquadric $Q(\infty, B)$, where $B \in \mathbb{N}_0 \cup \{\infty\}$; see Section 7. Lempert [15] shows that any strictly pseudoconvex real-analytic compact hypersurface can be mapped into a sphere in a possibly infinite dimensional Hilbert space. We show that if $Q(a, b)$ is not equivalent to a sphere, then for a CR mapping $Q(a, b) \to Q(\infty, B)$ where the image is not contained in a hyperplane we must have $B = \infty$. However, we also construct mappings whose image is not contained in a hyperplane from an $M \subset \mathbb{C}^n$ with an indefinite nondegenerate Levi-form to $Q(\infty, B)$ for finite B.

The authors would like to acknowledge John D'Angelo for many useful conversations on the subject and suggestions relating to this project. The second author would also like to thank Peter Ebenfelt for useful discussions on this subject. Finally the authors would like to acknowledge MSRI and AIM for holding workshops on the subject of CR complexity, which the authors attended and which led to the present project.

2. The basic setup

We begin by recalling the relevant definitions and proving some basic results about restrictions of real-valued polynomials. Let $\langle \cdot, \cdot \rangle$ denote the standard Hermitian inner product. Let A be a Hermitian symmetric matrix, then we call $\langle Az, z \rangle$ a *Hermitian form* where $z \in \mathbb{C}^{n+1}$.

Let

$$\mathcal{Z}_{n,d} \colon \mathbb{C}^{n+1} \to \mathbb{C}^{\binom{n+d}{d}} \tag{11}$$

be the mapping whose components are all the degree d monomials in $n + 1$ variables $z_0, z_1, \ldots, z_n$. That is, $\mathcal{Z}_{n,d}$ is the mapping given by

$$z \mapsto (z_0^d, z_0^{d-1} z_1, \ldots, z_n^d). \tag{12}$$

The mapping $\mathcal{Z}_{n,d}$ is called the *Veronese mapping in $n + 1$ variables of degree d.*

Definition 2.1. Let L be an $(m + 1)$-dimensional subspace of $\mathbb{C}^{n+1}$. Let E_L be a linear embedding of L into $\mathbb{C}^{n+1}$. Define $S^d_{E_L}$ to be the linear mapping that makes the following diagram commute:

$$\begin{tikzcd} \mathbb{C}^{m+1} \arrow[r, "E_L"] \arrow[d, "\mathcal{Z}_{m,d}"] & \mathbb{C}^{n+1} \arrow[d, "\mathcal{Z}_{n,d}"] \\ \mathbb{C}^{\binom{m+d}{d}} \arrow[r, "S^d_{E_L}"] & \mathbb{C}^{\binom{n+d}{d}}. \end{tikzcd} \tag{13}$$

Define $T^d_{E_L}$ to be the matrix associated to the linear mapping $S^d_{E_L}$, then

$$\mathcal{Z}_{n,d} \circ E_L = T^d_{E_L} \mathcal{Z}_{m,d} \tag{14}$$

We often abbreviate T^d_L for $T^d_{E_L}$ when the particular choice of embedding is not crucial.

Note that $T^d_{E_L}$ is not unique; it depends on the particular embedding E_L. However if we have E_L and F_L two embeddings of L then we have $T^d_{E_L} = \psi \circ T^d_{F_L}$ where ψ is an isomorphism.

Lemma 2.2. *Let $r(z, \bar{z})$ be a bihomogeneous polynomial defined for $z \in \mathbb{C}^{n+1}$. If ϕ is a linear change of coordinates of $\mathbb{C}^{n+1}$, then*

$$\operatorname{rank} r \circ \phi = \operatorname{rank} r. \tag{15}$$

Proof. We prove a more general fact in Lemma 5.2. ☐

The lemma shows that rank of $r|_{E_L}$ does not depend on the choice of embedding E_L. We state this observation in the following lemma.

Lemma 2.3. *Let $r(z, \bar{z}) = \langle C\mathcal{Z}_{n,d}, \mathcal{Z}_{n,d} \rangle$ where C is a Hermitian matrix, and suppose $n \geq 1$. Let L be an $(m+1)$-dimensional subspace of $\mathbb{C}^{n+1}$. Let E_L and F_L be two embeddings of L into $\mathbb{C}^{n+1}$. Then*

$$\operatorname{rank} T^*_{E_L} C T_{E_L} = \operatorname{rank} T^*_{F_L} C T_{F_L}. \tag{16}$$

Definition 2.4. Let $H_{\vec{c}}$ denote the hyperplane in $\mathbb{C}^{n+1}$ given by the following defining equation:

$$z_0 = c_1 z_1 + \cdots + c_n z_n, \tag{17}$$

where $\vec{c} = (c_1, c_2, \ldots, c_n)$. We embed $H_{\vec{c}}$ using $E_{H_{\vec{c}}}(z_1, \ldots, z_n) = (\sum_{i=1}^n c_i z_i, z_1, \ldots, z_n)$.

We abbreviate the notation for the restriction matrix $T^d_{H_{\vec{c}}}$ by $T^d_{\vec{c}}$.

The restriction matrix $T^d_{\vec{c}}$ is of size $\binom{n+d}{d} \times \binom{n+d-1}{d}$.

To study the maximum rank of restrictions of bihomogeneous polynomials to hyperplanes, it suffices to study the hyperplanes of the form $H_{\vec{c}}$.

Lemma 2.5. *Let $r(z, \bar{z})$ be a bihomogeneous polynomial defined for $z \in \mathbb{C}^{n+1}$ and suppose $n \geq 1$. Then*

$$\max_H \operatorname{rank} r|_H = \max_{\vec{c} \in \mathbb{C}^n} \operatorname{rank} r|_{H_{\vec{c}}}, \tag{18}$$

where H ranges over all complex hyperplanes through the origin.

Proof. The set of hyperplanes of the form $H_{\vec{c}}$ is dense in the Grassmannian $G(n, n+1)$. The rank of the restriction being bounded by a fixed integer is a holomorphic condition. Therefore $\operatorname{rank} r|_H$ achieves the maximum on an open dense set in $G(n, n+1)$, and the conclusion follows. ☐

3. Green's Restriction Theorem

The main technical tool in this paper is Green's hyperplane restriction theorem from [12]. We begin this section by recalling Macaulay representations and their basic properties. Then we give a precise statement of Green's theorem and translate the result into the language of this paper.

Given a non-negative integer c and a positive integer d, then c can be uniquely expressed in the following form:

$$c = \binom{k_d}{d} + \binom{k_{d-1}}{d-1} + \cdots + \binom{k_1}{1}, \tag{19}$$

where $0 \leq k_1 < k_2 < \cdots < k_d$. This expression is known as the d-th Macaulay representation of c. Following [12], we introduce the following notation:

$$c_{<d>} = \binom{k_d - 1}{d} + \binom{k_{d-1} - 1}{d-1} + \cdots + \binom{k_1 - 1}{1} \tag{20}$$

where we define $\binom{a}{b} = 0$ for $a < b$. Notice that $\cdot_{<d>}$ is monotone increasing.

Theorem 3.1 (Green [12]). *Let E be a linear system in $H^0(\mathbb{CP}^n, \mathcal{O}(d))$ and E_H be the restriction of E to a general hyperplane $H \in G(n, n+1)$. If c is the codimension of E in $H^0(\mathbb{CP}^n, \mathcal{O}(d))$ and c_H is the codimension of E_H in $H^0(\mathbb{CP}^{n-1}, \mathcal{O}(d))$, then*

$$c_H \leq c_{<d>}. \tag{21}$$

We seek a bound of the dimension of E in terms of the generic dimension of the restriction. Thus we restate the result in this way. First we restate the theorem in terms of matrices. If E is a linear system in $H^0(\mathbb{CP}^n, \mathcal{O}(d))$ then we can write: $E = \operatorname{span}\{s_1, s_2, \ldots, s_m\}$ where $s_i \in H^0(\mathbb{CP}^n, \mathcal{O}(d))$. Since each s_i is a degree d homogeneous polynomial in $z_0, z_1, \ldots, z_n$, then we can rewrite as $s_i = A_i \mathcal{Z}_{n,d}$ where we think of each A_i as a vector of the right dimension. Then we have $\operatorname{rank} E = \operatorname{rank} A$ where A is the matrix whose rows are A_i.

If we restrict E to a hyperplane H we obtain

$$E|_H = \operatorname{span}\{s_1|_H, s_2|_H, \ldots, s_m|_H\}. \tag{22}$$

Then $s_i|_{H_{\vec{c}}} = A_i T^d_{\vec{c}} \mathcal{Z}_{n-1,d}$, and therefore

$$\operatorname{rank} E|_{H_{\vec{c}}} = \operatorname{rank} A T^d_{\vec{c}}. \tag{23}$$

If we refer to a generic choice of H then equivalently, we refer to a generic set of c_i's, where we have: $\operatorname{rank}_{\mathbb{C}} E|_{H_{\vec{c}}} = \operatorname{rank}_{\mathbb{C}[c_1,\ldots,c_n]} A T^d_{\vec{c}}$. That is, the rank of $E|_{H_{\vec{c}}}$ over $\mathbb{C}$ is equal to the rank of $A T^d_{\vec{c}}$ over the polynomial ring $\mathbb{C}[c_1, \ldots, c_n]$.

Green's theorem can now be restated as follows: Let A be any matrix of size $m \times \binom{n+d}{d}$, and $N = \operatorname{rank}_{\mathbb{C}} A$. Recall the matrix $T^d_{\vec{c}}$ is of size $\binom{n+d}{d} \times \binom{n-1+d}{d}$. Let $k = \operatorname{rank}_{\mathbb{C}[c_1,\ldots,c_n]} A T^d_{\vec{c}}$. Then we have the following relationship:

$$G(n, d, N) \leq k \tag{24}$$

where

$$G(n, d, N) = \binom{n+d-1}{d} - \left(\binom{n+d}{d} - N \right)_{<d>}. \tag{25}$$

In the following lemma, we show that the lower bound G is constant in degree.

Lemma 3.2. *If $N \leq \binom{n+d}{d}$, $d \geq 1$, and $n \geq 2$, then $G(n, d, N) = G(n, d+1, N)$.*

Proof. The lemma follows immediately from

$$\left[\binom{n+d+1}{d+1}-N\right]_{<d+1>}=\binom{n+d-1}{d+1}+\left[\binom{n+d}{d}-N\right]_{<d>}. \tag{26}$$

□

By the lemma, we know that G is constant in degree when it is defined. We also have G is monotone in N and $G(n,d,\binom{n+d}{d})=\binom{n+d-1}{d}$, hence the following bound is finite and well-defined for fixed k.

Definition 3.3. Fix n and k. Define

$$K_n(k):=\max\left\{N_0\in\mathbb{N}_0 : G(n,d,N_0)\le k,\ \text{where } d \text{ is such that } N_0\le\binom{n+d}{d}\right\}. \tag{27}$$

Remark 3.4. The function $K_n(k)$ gives the sharp dimension bound from Green's theorem. The bound is achieved by restricting all the monomials in n variables of degree d. In two dimensions, $K_2(k)$ has a particularly simple form:

$$K_2(k)=\frac{k(k+1)}{2}. \tag{28}$$

One can show that $K_n(k)\le\frac{k(k+1)}{2}$, but this bound is not sharp in general.

The version of Green's theorem that we use in this paper is recorded in the following lemma. The important point is that the bound $K_n(k)$ does not depend on the degree d.

Lemma 3.5. *Let $n\ge 2$. If $A\mathcal{Z}_{n,d}$ is a linear system in $H^0(\mathbb{CP}^n,\mathcal{O}(d))$ with rank N and k is the rank of the restriction to a general hyperplane; namely, $k=\operatorname{rank}_{\mathbb{C}[c_1,\dots,c_n]} AT^d_{\vec{c}}$, then*

$$N\le K_n(k). \tag{29}$$

The lemma gives an upper bound for N that is independent of d for fixed k and $n\ge 2$. No such upper bound is possible if $n=1$.

4. Restrictions of Hermitian forms

In this section, we prove Theorem 1.2. To illustrate the main idea of the proof we prove a weaker version of the main result for hyperplanes.

Lemma 4.1. *Let $r(z,\bar z)$ be a real-valued bihomogeneous polynomial defined for $z\in\mathbb{C}^{n+1}$, and suppose $n\ge 2$. Then there exists a function $R_n\colon\mathbb{N}_0\to\mathbb{N}_0$ such that*

$$\operatorname{rank} r\le R_n\left(\max_H \operatorname{rank} r|_H\right), \tag{30}$$

where H ranges over all complex hyperplanes through the origin.

Proof. Suppose r is a nonzero bihomogeneous polynomial of bidegree (d,d). We write $r(z,\bar z)=\langle C\mathcal{Z}_{n,d},\mathcal{Z}_{n,d}\rangle$, where C is the coefficient matrix. Restricting r to a hyperplane H, we have

$$r|_H(z,\bar z)=\langle T_H^*CT_H\mathcal{Z}_{n-1,d},\mathcal{Z}_{n-1,d}\rangle. \tag{31}$$

Let $k=\max_H \operatorname{rank} r|_H=\max_H\operatorname{rank} T_H^*CT_H$ where H ranges over all complex hyperplanes, then by Lemma 2.5,

$$k=\max_{\vec{c}\in\mathbb{C}^n}\operatorname{rank} r|_{H_{\vec{c}}}=\max_{\vec{c}}\operatorname{rank}\,(T^d_{\vec{c}})^*CT^d_{\vec{c}}\ge\operatorname{rank}_{\mathbb{C}[\bar c_1,\dots,\bar c_n,c_1,\dots,c_n]}(T^d_{\vec{c}})^*CT^d_{\vec{c}}. \tag{32}$$

The strategy is to polarize and use Lemma 3.5 twice.

Let $D = CT^d_{\vec{c}}$, $m = \operatorname{rank}_{\mathbb{C}} C$, and let $k_1 = \operatorname{rank}_{\mathbb{C}[c_1,\ldots,c_n]} D$. By Lemma 3.5, we have

$$m \leq K_n(k_1). \tag{33}$$

For a generic choice of $c_1, \ldots, c_n$, we have

$$k_1 = \operatorname{rank}_{\mathbb{C}} D = \operatorname{rank}_{\mathbb{C}} D^*. \tag{34}$$

Likewise, polarizing and taking generic choices of $\bar{c}_1, \ldots, \bar{c}_n$, we have

$$\operatorname{rank}_{\mathbb{C}[\bar{c}_1,\ldots,\bar{c}_n,c_1,\ldots,c_n]} (T^d_{\vec{c}})^* CT^d_{\vec{c}} = \operatorname{rank}_{\mathbb{C}[c_1,\ldots,c_n]} D^* T^d_{\vec{c}} \tag{35}$$

Applying Lemma 3.5 again, we get

$$k_1 \leq K_n(k). \tag{36}$$

Thus $\operatorname{rank} r \leq K_n(K_n(k))$, and we have an upper bound for $\operatorname{rank} r$ that is independent of degree. We complete the proof by defining $R_n(k)$ to be the maximum $\operatorname{rank} r$ for any bihomogeneous polynomial r with $\max_H \operatorname{rank} r|_H \leq k$. ☐

Remark 4.2. In the proof of Lemma 4.1, we proved that

$$R_n(k) \leq K_n\bigl(K_n(k)\bigr). \tag{37}$$

In fact, we believe that $R_n(k) = K_n(k)$.

We are now ready to prove the main result, Theorem 1.2. For reader convenience we restate the theorem.

Theorem. *Let $n \geq 2$, $z \in \mathbb{C}^{n+1}$, and $1 \leq m \leq n-1$. Then there exists a function $R_{m,n} \colon \mathbb{N}_0 \to \mathbb{N}_0$ such that for any bihomogeneous polynomial $r(z,\bar{z})$,*

$$\operatorname{rank} r \leq R_{m,n}\left(\max_{L \in G(m+1,n+1)} \operatorname{rank} r|_L \right). \tag{38}$$

Proof. Suppose r is a nonzero bihomogeneous polynomial of bidegree (d,d). We again write $r(z,\bar{z}) = \langle C\mathcal{Z}_{n,d}, \mathcal{Z}_{n,d}\rangle$. After applying a linear change of coordinates we can assume without loss of generality that all $L \in G(m+1,n+1)$ are given as

$$z' = Vz'', \tag{39}$$

for $(z',z'') \in \mathbb{C}^{n-m} \times \mathbb{C}^{m+1}$ and $V \in M_{n-m,m+1}$ is a matrix. Using the embedding $z'' \mapsto (Vz'', z'')$ we obtain a restriction matrix T_L. We notice that T_L depends holomorphically on the entries of V.

We write

$$r|_L(z,\bar{z}) = \langle T_L^* C T_L \mathcal{Z}_{m,d}, \mathcal{Z}_{m,d}\rangle. \tag{40}$$

And again note that

$$\operatorname{rank} r|_L = \operatorname{rank} T_L^* C T_L \tag{41}$$

If $m = n-1$ we are done. We let $R_{n-1,n}(k) = R_n(k)$. Otherwise, T_L is a composition $T_L = T_S T_H$ where H is a hyperplane through the origin in S, where S is an $m+2$ dimensional plane through the origin in $\mathbb{C}^{n+1}$. As $T_L^* C T_L = (T_H T_S)^* C T_S T_H$ is of rank less than k, $T_S^* C T_S$ is of rank less than $R_m(k)$ by Lemma 4.1. The theorem follows by applying Lemma 4.1 $n-m$ times. The function $R_{m,n}(k)$ is then defined recursively by composition. ☐

We say a set $\mathcal{L} \subset G(m+1,n+1)$ is *generic* if $\mathcal{L}$ is not contained in any proper complex algebraic subvariety of $G(m+1,n+1)$. That is, if no nontrivial polynomial vanishes on $\mathcal{L}$. With this terminology, we have the following corollary:

Corollary 4.3. *Let $r(z,\bar{z})$ be a positive semidefinite bihomogeneous polynomial defined for $z \in \mathbb{C}^{n+1}$, $n \geq 2$. Let $1 \leq m \leq n-1$. Then there exists a function $K_{m,n} \colon \mathbb{N}_0 \to \mathbb{N}_0$ such that*

$$\operatorname{rank} r \leq K_{m,n}\left(\max_{L\in\mathcal{L}} \operatorname{rank} r|_L\right). \tag{42}$$

where $\mathcal{L} \subset G(m+1,n+1)$ is a generic subset.

Note that $r(z,\bar{z}) \geq 0$ for all $z \in \mathbb{C}^{n+1}$ is not enough to conclude that the matrix of coefficients is positive semidefinite. On the other hand, if the matrix is positive semidefinite, then $r(z,\bar{z}) \geq 0$ is true for all $z \in \mathbb{C}^{n+1}$. See [8,9] for more on positivity conditions.

Proof. The proof as before carries on after we make the following observation. Let $r(z,\bar{z}) = \langle C\mathcal{Z}_{n,d}, \mathcal{Z}_{n,d}\rangle$. If C is positive semidefinite then we can write $C = A^*A$. In the restriction:

$$r|_L(z,\bar{z}) = \langle T_L^* C T_L \mathcal{Z}_{m,d}, \mathcal{Z}_{m,d}\rangle. \tag{43}$$

And again note that

$$\operatorname{rank} r|_L = \operatorname{rank} T_L^* C T_L = \operatorname{rank} AT_L. \tag{44}$$

Let $k = \max_{L\in\mathcal{L}} \operatorname{rank} r|_L$. The condition

$$\operatorname{rank} AT_L \leq k \tag{45}$$

is defined by the vanishing of certain polynomials in the entries of AT_L. Defining L with a matrix $V \in M_{n-m,m+1}$ as in (39), we notice that as A is fixed, then (45) is defined by vanishing of certain polynomials in the entries of V. We know that (45) is true for all $L \in \mathcal{L}$, where $\mathcal{L}$ is not contained in any complex algebraic subvariety of $G(m+1,n+1)$. The set of corresponding matrices V is not contained in any complex algebraic subvariety of $M_{n-m,m+1}$. Thus (45) holds for all $V \in M_{n-m,m+1}$, and so for all $L \in G(m+1,n+1)$.

The conclusion follows by applying Lemma 3.5 $n-m$ times. The function $K_{m,n}(k)$ is then defined recursively by composition as in the proof of Theorem 1.2. ☐

5. Real-analytic case

We wish to decompose real-analytic functions as a difference of squared norms of Hilbert space valued holomorphic functions. This line of reasoning follows the ideas pioneered by D'Angelo [5], although we introduce a rescaling that allows us to work with bounded operators.

First let us prove that the matrix of coefficients is a trace-class operator if a series converges in the right neighborhood. See the book [1] for more information on matrix operators on ℓ^2. Let $\mathcal{Z} = \mathcal{Z}_n = (\ldots, z^\alpha, \ldots)$ be the mapping of $z \in \mathbb{C}^n$ to the space of infinite sequences, where components of the mapping are all possible monomials. By using the geometric series we note that $\mathcal{Z}$ maps the unit polydisc Δ to ℓ^2.

Lemma 5.1. *Let $\Delta \subset \mathbb{C}^n$ be the unit polydisc. Suppose that $r(z,\bar{z})$ is a real-analytic function whose complexified power series converges in a neighborhood of $\overline{\Delta} \times \overline{\Delta} \subset \mathbb{C}^n \times \mathbb{C}^n$. Then the matrix of coefficients of r defines a trace-class operator on ℓ^2.*

Proof. Write r as $\langle C\mathcal{Z}, \mathcal{Z}\rangle$ where $C = [c_{\alpha\beta}]_{\alpha\beta}$ is the matrix of coefficients. The hypothesis says that the interior of the domain of convergence includes the point $z = (1,1,\ldots,1)$. As the series converges absolutely at $(1,1,\ldots,1)$ we get $\sum_{\alpha\beta} |c_{\alpha\beta}| < \infty$. A matrix whose entries are summable defines a compact operator. Moreover, if $\{e_\alpha\}$ is the standard basis corresponding to the monomials, then

$$\sum_\alpha \langle |C|\, e_\alpha, e_\alpha\rangle \leq \sum_\alpha \bigl\| |C|\, e_\alpha \bigr\| = \sum_\alpha \| C e_\alpha \| \leq \sum_\alpha \sum_\beta \left| \langle C e_\alpha, e_\beta \rangle \right| < \infty. \tag{46}$$

Here $|C|$ is the unique positive Hermitian square root of C^*C. So C is trace-class. ☐

As C is a Hermitian trace-class operator, we apply the spectral theorem. We write C as a sum of rank one matrices as follows. Let λ_k be the k-th nonzero eigenvalue. Let $\epsilon_k = \pm 1$ be the sign of the k-th nonzero eigenvalue. We ignore all zero eigenvalues. Let $\{v_k\}$ be an orthonormal set of corresponding eigenvectors. Let $m \in \mathbb{N}_0 \cup \{\infty\}$ be the rank of C, then we write

$$C = \sum_{j=1}^{m} \lambda_j v_j v_j^*. \tag{47}$$

The sum converges as the eigenvalues of a trace-class operator are absolutely summable. We define holomorphic functions $f_j(z) = \sqrt{|\lambda_j|}\, v_j^* \mathcal{Z}$. As $v_j \in \ell^2$, we note that each f_j converges in the unit polydisc. We see that

$$\langle C\mathcal{Z}, \mathcal{Z} \rangle = \sum_{j=1}^{m} \epsilon_j \left| f_j(z) \right|^2. \tag{48}$$

Suppose that for some m', $\epsilon'_j = \pm 1$, and some g_j holomorphic in the unit polydisc we write

$$\langle C\mathcal{Z}, \mathcal{Z} \rangle = \sum_{j=1}^{m'} \epsilon'_j \left| g_j(z) \right|^2. \tag{49}$$

If $m' = \infty$, then obviously $m' \geq m$. If m' is finite then we have written C as a sum of m' rank-one operators, and hence $m' \geq m$.

Next we have to show that the rank of the matrix C is unchanged under changes of coordinates. We define the rank of r at the origin to be the rank of the matrix C.

Lemma 5.2. *Let $\Delta \subset \mathbb{C}^n$ be the unit polydisc. Suppose that $r(z,\bar{z})$ is a real-analytic function whose complexified power series converges in a neighborhood of $\overline{\Delta} \times \overline{\Delta} \subset \mathbb{C}^n \times \mathbb{C}^n$.*

Suppose that φ is a biholomorphic change of coordinates that takes a neighborhood of $\overline{\Delta}$ to a neighborhood of $\overline{\Delta}$. Then $r \circ \varphi$ (which also converges on a neighborhood of $\overline{\Delta} \times \overline{\Delta} \subset \mathbb{C}^n \times \mathbb{C}^n$) has the same rank as r at the origin.

Proof. Suppose that r is of finite rank k. Then there exist k eigenvectors of C and therefore we can write

$$r(z,\bar{z}) = \sum_{j=1}^{k} \epsilon_j \left| f_j(z) \right|^2 \tag{50}$$

for $\epsilon_j = \pm 1$ and f_j holomorphic functions. Therefore

$$(r \circ \varphi)(z,\bar{z}) = \sum_{j=1}^{k} \epsilon_j \left| f_j(\varphi(z)) \right|^2, \tag{51}$$

so $\operatorname{rank} r \circ \varphi \leq \operatorname{rank} r$. We are finished by symmetry. ☐

We can also define the number of positive and negative eigenvalues at a point (the rank of the positive or negative definite part). Using exactly the same argument as above we can show that the rank, the number of positive and the number of negative eigenvalues are constant on connected sets.

Lemma 5.3. *Let $U \subset \mathbb{C}^n$ be a connected open set. Let $r \colon U \to \mathbb{R}$ be a real-analytic function. Let $p_1, p_2 \in U$ and let k_j, a_j, b_j denote the rank, the number of positive, and the number of negative eigenvalues respectively at p_j. Then $k_1 = k_2$, $a_1 = a_2$, and $b_1 = b_2$.*

With the aid of the lemmas above we can define the rank of r regardless of where it converges. We simply translate and dilate r so that it converges in the unit polydisc and then take the rank of the resulting compact operator.

If the domain U of r is connected, we define the *rank* of r as the rank at some fixed point of U. Similarly define the *signature pair* to be the pair (a, b) if r has a positive and b negative eigenvalues. Of course we allow a and b be in $\mathbb{N}_0 \cup \{\infty\}$.

Lemma 5.4. *Let L be an affine complex submanifold of $\mathbb{C}^n$ of dimension m that intersects the unit polydisc $\Delta_n \subset \mathbb{C}^n$ and $E \colon \mathbb{C}^m \to L$ an embedding such that E takes the closed unit polydisc $\overline{\Delta}_m \subset \mathbb{C}^m$ to a subset of $L \cap \Delta_n$.*

The restriction matrix T_L defined by $\mathcal{Z}_n \circ E = T_L \mathcal{Z}_m$ is a Hilbert-Schmidt operator.

Proof. Note that $\|\mathcal{Z}_n\|^2 = \langle \mathcal{Z}_n, \mathcal{Z}_n \rangle$ is the absolute value squared of the geometric series and hence converges in the unit polydisc $\Delta_n \subset \mathbb{C}^n$. As E takes the closed unit polydisc $\overline{\Delta}_m \subset \mathbb{C}^m$ to Δ_n, we have that $\|\mathcal{Z}_n \circ E\|^2$ converges absolutely for $z \in \overline{\Delta}_m$, therefore

$$\|\mathcal{Z}_n \circ E\|^2 = \langle \mathcal{Z}_n \circ E, \mathcal{Z}_n \circ E \rangle = \langle T_L^* T_L \mathcal{Z}_m, \mathcal{Z}_m \rangle, \tag{52}$$

where T_L is the restriction matrix induced by the embedding E. By Lemma 5.1, $T_L^* T_L$ is trace-class and so T_L is a Hilbert-Schmidt operator. ☐

When $r(z, \bar{z}) = \langle C\mathcal{Z}, \mathcal{Z} \rangle$ is positive semi-definite, that is, has no negative eigenvalues, we can always write

$$r(z, \bar{z}) = \langle A^* A \mathcal{Z}, \mathcal{Z} \rangle = \|A\mathcal{Z}\|^2 .$$

When everything is scaled appropriately so that C is trace-class, then A is Hilbert-Schmidt.

We now prove the real-analytic version of the main theorem on rank of Hermitian forms for positive semi-definite forms. The proof reduces to the algebraic version. However, note that a key point here is that the rank of r is not required to be finite. As before a generic subset $\mathcal{L} \subset G_{m,n}$ is a subset not contained in any proper complex algebraic subvariety of $G_{m,n}$, that is, no nontrivial polynomial vanishes on $\mathcal{L}$.

Lemma 5.5. *Let $r(z, \bar{z})$ be a positive semi-definite real-analytic function whose complexified power series converges in a neighborhood of $\overline{\Delta} \times \overline{\Delta} \subset \mathbb{C}^n \times \mathbb{C}^n$. Let $1 \leq m \leq n-1$, and let $\mathcal{L} \subset G_{m,n}$ be a generic subset such that all the corresponding manifolds intersect Δ. Further suppose that*

$$\sup_{L \in \mathcal{L}} \operatorname{rank} r|_L < \infty. \tag{53}$$

Then $\operatorname{rank} r$ *is finite and moreover there exists a function $K_{m,n} \colon \mathbb{N}_0 \to \mathbb{N}_0$ such that*

$$\operatorname{rank} r \leq K_{m,n} \left(\max_{L \in \mathcal{L}} \operatorname{rank} r|_L \right). \tag{54}$$

Proof. For an affine manifold L that passes through the unit polydisc, we can find an embedding $E\colon \mathbb{C}^m \to H$ such that E takes the closed unit polydisc $\overline{\Delta}_m \subset \mathbb{C}^m$ to a subset of $H \cap \Delta_n$. The induced restriction matrix T_L is a Hilbert-Schmidt operator by Lemma 5.4.

As before, we assume that L is a graph over the first m variables and we assume that it contains a point of the set $\{z\colon z_1 = z_2 = \cdots = z_m = 0, |z_j| < 1 \text{ for } j = m+1,\ldots,n\}$. So assume that $\mathcal{L}$ only contains such L. As in the proof of Theorem 1.2 we identify $\mathcal{L}$ with a subset $\mathcal{V} \subset M_{n-m,m+1}$. For such L we can simply take a dilation of the first m variables to obtain an embedding E that maps the closed unit polydisc $\overline{\Delta}_m$ into Δ_n, to ensure that T_L is a Hilbert-Schmidt operator. Without loss of generality, we can also assume that the dilation is the same for all $L \in \mathcal{L}$. Therefore we can again assume that entries of T_L depend holomorphically on the entries of $V \in \mathcal{V}$.

Let $\alpha = \max_{L\in\mathcal{L}} \operatorname{rank} r|_L$. Write

$$r(z,\bar{z}) = \langle A^*A\mathcal{Z}, \mathcal{Z}\rangle. \tag{55}$$

Thus for $L \in \mathcal{L}$ we have

$$\operatorname{rank} AT_L \leq \alpha. \tag{56}$$

As for the finite case, this condition is defined by the vanishing of polynomials in the entries of AT_L, these depend holomorphically on the entries of V. Thus (56) is true for all affine manifolds of dimension m that intersect Δ.

Unless $m = n-1$ we again write $T_L = T_S T_H$ for a hyperplane H in S, where S is an affine manifold of dimension $m+1$. Hence if we prove the result of the theorem for hyperplanes, the general case follows by induction as before.

Therefore assume that

$$\operatorname{rank} r|_H \leq \alpha. \tag{57}$$

for all hyperplanes H that intersect Δ. Then we have that $\operatorname{rank} AT_H \leq \alpha$ for all such hyperplanes H.

Let us suppose that $H = H_{\vec{c}}$ is the hyperplane given by the vector $\vec{c}$ as before. We take T_H and look at the submatrix of rows and columns corresponding to monomials of degree d or less, for some integer d. Then we see that

$$T_H = \begin{bmatrix} T'^{d}_{\vec{c}} & * \\ 0 & * \end{bmatrix}, \tag{58}$$

where $T'^{d}_{\vec{c}} = T^{d}_{\vec{c}} D$, where $T^{d}_{\vec{c}}$ is given by Definition 2.4 and D is an invertible diagonal matrix (coming from the dilation used above).

Let us first suppose that A is of finite rank N. We can therefore assume that A is a $N \times \infty$ matrix. We have that $\operatorname{rank} AT_H \leq \alpha$ for all H. Write $A = [A' \ A'']$, then

$$AT_H = [A'T'^{d}_{\vec{c}} \quad *]. \tag{59}$$

Therefore, $\operatorname{rank} A'T'^{d}_{\vec{c}} = \operatorname{rank} A'T^{d}_{\vec{c}} \leq \alpha$ for all $\vec{c}$. As A is of finite rank then for some large d, A' contains enough columns such that $\operatorname{rank} A' = N$.

We have therefore reduced the proof of the analytic case to the same matrix $T^{d}_{\vec{c}}$ and to finish the proof for A of finite rank N, it is enough to apply Lemma 3.5. That is, the rank of A and therefore r are bounded by $K_n(\alpha)$.

Now let us drop the assumption that the rank of A is finite. Let A_ν be an infinite by infinite matrix consisting of the first ν rows of A followed by all zero rows. Obviously A_ν is of finite rank. We have that

$$\operatorname{rank} A_\nu T_H \leq \operatorname{rank} AT_H \leq \alpha. \tag{60}$$

It follows that $\operatorname{rank} A_\nu \leq K_n(\alpha)$ for all ν.

From the proof of Lemma 5.1 we note that the entries of A^*A are absolutely summable and hence the entries of A are square summable. Therefore A_ν converges to A in the Hilbert-Schmidt norm. Then as the rank of A_ν is uniformly bounded by $K_n(\alpha)$, we obtain

$$\operatorname{rank} A \leq K_n(\alpha). \tag{61}$$

To see this fact, simply pick a finite orthonormal set $\{Aw_j\}$ in the range of A and then notice that $A_\nu w_j$ must converge to Aw_j as $\nu \to \infty$. Thus for large enough ν, the set $\{A_\nu w_j\}$ must be linearly independent (and hence of cardinality less than $K_n(\alpha)$). ☐

Example 5.6. Let us show that positivity of r is necessary. First note that the set $\mathcal{L}$ of affine complex manifolds of dimension 1 that lie in

$$Q(2,1) = \{z \in \mathbb{C}^3 : |z_1|^2 + |z_2|^2 - |z_3|^2 = 1\} \tag{62}$$

forms a generic set in $G_{1,3}$. See Section 6 for a proof of this fact. Let

$$r(z,\bar{z}) = (|z_1|^2 + |z_2|^2 - |z_3|^2)^d. \tag{63}$$

Then for $z \in Q(2,1)$

$$r(z,\bar{z}) = (|z_1|^2 + |z_2|^2 - |z_3|^2)^d = 1. \tag{64}$$

Therefore the rank of $r|_L$ for $L \in \mathcal{L}$ is always 1. However the rank of r is $\binom{2+d}{d}$.

By adding a function of infinite rank to r that vanishes on $Q(2,1)$ we can obtain an example of r with rank of $r|_L$ bounded but rank of r infinite.

We can now prove Theorem 1.1. That is, we can drop positivity of r if we assume that the rank is bounded on all hyperplanes. The proof is very similar to the proof of Lemma 4.1.

Proof of Theorem 1.1. Suppose r is a nonzero Hermitian form. We can assume that r converges in a neighbourhood of the closed unit polydisc as above. Write $r(z,\bar{z}) = \langle C\mathcal{Z}_n, \mathcal{Z}_n\rangle$. It is again enough to prove the theorem for hyperplanes. Restricting r to a hyperplane H that intersects the closed unit polydisc, we have

$$r|_H(z,\bar{z}) = \langle T_H^* C T_H \mathcal{Z}_{n-1}, \mathcal{Z}_{n-1}\rangle. \tag{65}$$

Let $k = \max_H \operatorname{rank} r|_H = \max_H \operatorname{rank} T_H^* C T_H$ where H ranges over all complex hyperplanes. As before, the set of hyperplanes $H_{\vec{c}}$ is open and dense in the set of all hyperplanes, so

$$k = \max_{\vec{c} \in \mathbb{C}^n} \operatorname{rank} r|_{H_{\vec{c}}} = \max_{\vec{c}} \operatorname{rank} (T_{\vec{c}}^d)^* C T_{\vec{c}}^d \geq \operatorname{rank}_{\mathbb{C}[\bar{c}_1,\dots,\bar{c}_n,c_1,\dots,c_n]} (T_{\vec{c}}^d)^* C T_{\vec{c}}^d. \tag{66}$$

Again we polarize and apply the semi-definite result twice.

Let $D = CT_{\vec{c}}^d$, $m = \operatorname{rank}_{\mathbb{C}} C$, and let $k_1 = \operatorname{rank}_{\mathbb{C}[c_1,\dots,c_n]} D$. Note that we still allow the ranks to be infinite. Applying the bound (61) from the proof of Lemma 5.5 we obtain

$$m \leq K_n(k_1) \tag{67}$$

For a generic choice of $c_1, \dots, c_n$, we have

$$k_1 = \operatorname{rank}_{\mathbb{C}} D = \operatorname{rank}_{\mathbb{C}} D^*. \tag{68}$$

Likewise, polarizing and taking generic choices of $\bar{c}_1, \dots, \bar{c}_n$, we have

$$\operatorname{rank}_{\mathbb{C}[\bar{c}_1,\dots,\bar{c}_n,c_1,\dots,c_n]} (T_{\vec{c}})^* C T_{\vec{c}} = \operatorname{rank}_{\mathbb{C}[c_1,\dots,c_n]} D^* T_{\vec{c}} \tag{69}$$

Again applying the bound (61) we get

$$k_1 \leq K_n(k). \tag{70}$$

Now we note that as k is finite, then k_1 is finite and hence $m \leq K_n\bigl(K_n(k)\bigr)$. Thus defining $R_n(k)$ as in the proof of Lemma 4.1 completes the proof. ☐

6. CR mappings of hyperquadrics

In this section we apply Theorem 1.1 on bounding the rank of Hermitian forms by the rank of restrictions to prove Theorem 1.3 on rigidity of CR mappings between hyperquadrics.

First let us note how to construct mappings from a real-analytic function $r(z,\bar{z})$ vanishing on the hyperquadric.

Proposition 6.1. *Let $p \in Q(a,b)$ be a point and let $\Omega \subset \mathbb{C}^{a+b}$ be a connected open set with $p \in \Omega$. Let $r \colon \Omega \to \mathbb{R}$ be nonzero real-analytic function of finite rank with A positive and B negative eigenvalues such that r vanishes on $Q(a,b) \cap \Omega$.*

Then there exists a point q arbitrarily close to p and a neighbourhood $U \subset Q(a,b)$ of q, and a real-analytic CR mapping $F \colon U \to Q(A,B-1)$ whose image is not contained in a hyperplane.

Proof. Write

$$r(z,\bar{z}) = \sum_{j=1}^{A} \left|f_j(z)\right|^2 - \sum_{j=1}^{B} \left|g_j(z)\right|^2, \tag{71}$$

with f_j and g_j linearly independent. The mapping

$$z \mapsto \Bigl(\frac{f_1(z)}{g_B(z)}, \dots \frac{f_A(z)}{g_B(z)}, \frac{g_1(z)}{g_B(z)}, \dots \frac{g_{B-1}(z)}{g_B(z)}\Bigr) \tag{72}$$

is a meromorphic mapping taking $Q(a,b)$ to $Q(A,B-1)$. If $g_B(p) \neq 0$ we are finished. Otherwise note that the set $\{g_B = 0\} \cap Q(a,b)$ is nowhere dense in $Q(a,b)$ and hence there is a point $q \in Q(a,b)$ arbitrarily close where $g_B(q) \neq 0$ and a holomorphic mapping defined near q taking $Q(a,b)$ to $Q(A,B-1)$. ☐

Therefore if r vanishes on $Q(a,b)$ and has signature pair (A,B) then r induces a mapping from $Q(a,b)$ to $Q(A,B-1)$.

Let us prove a lemma about signature pairs of functions vanishing on $Q(a,b)$ before proving Theorem 1.3.

Lemma 6.2. *Let $a \geq 2$, $b \geq 1$, and $a > b$. Let $\Omega \subset \mathbb{C}^{a+b}$ be a connected open set such that $\Omega \cap Q(a,b)$ is nonempty. Let $r \colon \Omega \to \mathbb{R}$ be a real-analytic function vanishing on $\Omega \cap Q(a,b)$ with signature pair (A,B). Then if $B < \infty$ we have*

$$A \leq M(a,b,B), \tag{73}$$

where $M = M(a,b,B)$ is a constant depending only a, b, and B.

Proof. Write

$$r(z,\bar{z}) = \sum_{j=1}^{A} \left|f_j(z)\right|^2 - \sum_{j=1}^{B} \left|g_j(z)\right|^2 = \left\lVert f(z)\right\rVert^2 - \left\lVert g(z)\right\rVert^2, \tag{74}$$

such that the components of f and g are linearly independent. Let us suppose that B is finite, but for now we allow f to map into an infinite dimensional Hilbert space.

After an affine change of coordinates we can assume that we are working in a neighborhood of the origin and that f and g converge in a neighborhood of a closed unit polydisc $\overline{\Delta}$. Let $\tilde{Q}(a,b)$ and $\tilde{\Omega}$ correspond to $Q(a,b)$ and Ω after this linear change of coordinates.

Let $\mathcal{Z}$ be the infinite vector of all monomials in z. We can then find an $A \times \infty$ matrix F such that $f(z) = F\mathcal{Z}$. Similarly we have a $B \times \infty$ matrix G such that $g(z) = G\mathcal{Z}$. We know that $\|f(z)\|^2$ converges in the closed unit polydisc, and so by Lemma 5.1 F defines a Hilbert-Schmidt operator (F^*F is trace-class) on ℓ^2. Similarly G also defines a Hilbert-Schmidt operator.

For $z \in \tilde{Q}(a,b) \cap \tilde{\Omega}$ we have

$$\|F\mathcal{Z}\|^2 - \|G\mathcal{Z}\|^2 = \|f(z)\|^2 - \|g(z)\|^2 = 0. \tag{75}$$

As f and g are linearly independent we have

$$\operatorname{rank} F = A \qquad \text{and} \qquad \operatorname{rank} G = B. \tag{76}$$

Let L be an affine complex submanifold of $\tilde{Q}(a,b)$. Taking the right embedding of L and applying Lemma 5.4, we find that the restriction matrix T_L is a Hilbert-Schmidt operator. Restricted to L we have that

$$\|F\mathcal{Z}\|^2 = \|G\mathcal{Z}\|^2 . \tag{77}$$

Thus have that $T_L F^* F T_L = T_L G^* G T_L$. In other words, rank of GT_L is equal to the rank of FT_L. The rank of GT_L is at most $B+1$. That means that the rank of FT_L is at most $B+1$ for all L that lie in $\tilde{Q}(a,b)$.

Let $(z', z'') \in \mathbb{C}^a \times \mathbb{C}^b$ be coordinates such that $Q(a,b)$ is defined by $\|z'\|^2 - \|z''\|^2 = 1$. We assume that $a > b$. The defining function for L that lie in $Q(a,b)$ is

$$z' = V \begin{bmatrix} z'' \\ 1 \end{bmatrix}, \tag{78}$$

where $V \in M_{a,b+1}$ (an $a \times (b+1)$ matrix) has orthonormal rows. Let us denote by L_V the affine manifold defined by such a V.

We claim that the set $\mathcal{V} \subset M_{a,b+1} \cong \mathbb{C}^{a(b+1)}$ of matrices with orthonormal columns is not contained in any proper complex algebraic subvariety of $\mathbb{C}^{a(b+1)}$. Suppose that there was a polynomial p in $a(b+1)$ variables that vanishes on $\mathcal{V}$. If we multiply the k-th column by $e^{i\theta_k}$ we note that p still has to vanish. By the uniqueness theorem for Fourier series we note that there has to be a polynomial that is independently homogeneous in entries of the k-th column (for all columns). That is, we can multiply each column independently by any complex number and still stay in the zero locus of p. The claim follows by working on each column independently now.

The claim shows that the set $\mathcal{L}$ of affine manifolds L corresponding to $V \in \mathcal{V}$ is a generic set in $G_{m,n}$. If we let $\tilde{\mathcal{L}} \subset G_{m,n}$ correspond to the set of affine manifolds that lie in $\tilde{Q}(a,b)$. We see that $\tilde{\mathcal{L}}$ must be generic as well.

We have all the ingredients to apply Lemma 5.5 to obtain that

$$A = \operatorname{rank} F \leq K_{b,a+b}(B). \tag{79}$$

□

Let us restate Theorem 1.3 for reader convenience before proving it.

Theorem. *Let $a > b \geq 1$, $U \subset Q(a,b)$ be a connected open set, and $f \colon U \to Q(A,B)$ be a real-analytic CR mapping such that $f(U)$ does not lie in a complex hyperplane then*

$$A \leq N(a,b,B), \tag{80}$$

where $N = N(a,b,B)$ is a constant depending only a, b, and B.

Proof. Let $\varphi = (f,g)$ denote a real-analytic CR mapping of $U \subset Q(a,b)$ to $Q(A,B)$, where $f \colon U \to \mathbb{C}^A$ and $g \colon U \to \mathbb{C}^B$. Plugging into the defining equation of the target hyperquadric we obtain that for $z \in U$

$$\|f(z)\|^2 - \|g(z)\|^2 - 1 = 0. \tag{81}$$

Real-analytic CR mappings extend to holomorphic mappings on a neighborhood of U. Therefore we can assume that f and g are holomorphic in a neighborhood of U. We let $r(z,\bar{z}) = \|f(z)\|^2 - \|g(z)\|^2 - 1$.

The condition that $\varphi(U)$ does not lie in a complex hyperplane is simply stating that components of the mapping $(f,g,1)$ are linearly independent. In particular we see that r has rank $A + B + 1$, and signature pair $(A, B+1)$.

We can now apply Lemma 6.2 to r to obtain

$$A \leq K_{b,a+b}(B+1). \tag{82}$$

□

7. Spheres and hyperquadrics in Hilbert space

Forstnerič [11] has shown that there exist strictly pseudoconvex real-analytic compact hypersurfaces that cannot be embedded into a sphere of any finite dimension. On the other hand, Lempert [15] has shown that a mapping exists if one takes the sphere in the infinite dimensional Hilbert space ℓ^2. Let us make the following natural definitions to extend hyperquadrics to ℓ^2:

$$Q(\infty,b) := \left\{ z \in \ell^2 : -\sum_{j=1}^{b} |z_j|^2 + \sum_{j=b+1}^{\infty} |z_j|^2 = 1 \right\}, \tag{83}$$

$$Q(\infty,\infty) := \left\{ z \in \ell^2 : \sum_{j=1}^{\infty} (|z_{2j-1}|^2 - |z_{2j}|^2) = 1 \right\}. \tag{84}$$

We write $S^\infty = Q(\infty,0)$ for the unit sphere in ℓ^2.

By a real-analytic CR mapping from a CR manifold M to $Q(\infty,b)$ we mean a holomorphic mapping to ℓ^2 defined on a neighborhood of M taking M to $Q(\infty,b)$.

We can use the ideas from the previous section to construct mappings. For example, consider the Hermitian form

$$r(z,\bar{z}) = \frac{e^{|z_1|^2+|z_2|^2} - 1}{e-1} - 1. \tag{85}$$

The function r vanishes precisely on the unit sphere in $\mathbb{C}^2$. In fact $\frac{e^{|z_1|^2+|z_2|^2}-1}{e-1}$ is 1 on the unit sphere and is positive semi-definite. Hence r has signature pair $(\infty,1)$; that is, r has infinitely positive eigenvalues and exactly one negative eigenvalue corresponding to the -1. Therefore the form induces a real-analytic CR mapping

$$f \colon S^2 \to S^\infty. \tag{86}$$

In particular we expand the series:

$$\begin{aligned}\frac{e^{|z_1|^2+|z_2|^2}-1}{e-1} &= \frac{(|z_1|^2+|z_2|^2)}{e-1}+\frac{(|z_1|^2+|z_2|^2)^2}{2(e-1)}+\frac{(|z_1|^2+|z_2|^2)^3}{6(e-1)}+\cdots \\ &= \left|\frac{z_1}{\sqrt{e-1}}\right|^2+\left|\frac{z_2}{\sqrt{e-1}}\right|^2+ \\ &\qquad +\left|\frac{z_1^2}{\sqrt{2(e-1)}}\right|^2+\left|\frac{z_1z_2}{\sqrt{2(e-1)}}\right|^2+\left|\frac{z_2^2}{\sqrt{2(e-1)}}\right|^2+\cdots\end{aligned} \tag{87}$$

Therefore

$$\begin{aligned}f(z_1,z_2,\bar z_1,\bar z_2) = \Big(&\frac{z_1}{\sqrt{e-1}},\frac{z_2}{\sqrt{e-1}},\frac{z_1^2}{\sqrt{2(e-1)}},\frac{z_1z_2}{\sqrt{2(e-1)}},\frac{z_2^2}{\sqrt{2(e-1)}}, \\ &\frac{z_1^3}{\sqrt{6(e-1)}},\frac{z_1^2z_2}{\sqrt{6(e-1)}},\frac{z_1z_2^2}{\sqrt{6(e-1)}},\frac{z_2^3}{\sqrt{6(e-1)}},\ldots\Big).\end{aligned} \tag{88}$$

Note that the image is not contained in a hyperplane; this fact can even be seen directly by uniqueness of power series.

By taking

$$r_2(z,\bar z) = r(z,\bar z)+|\varphi(z)|^2\, r(z,\bar z) \tag{89}$$

for an arbitrary holomorphic function $\varphi(z)$, we obtain a new form with signature pair $(\infty,2)$, and therefore a CR mapping $f\colon S^3\to Q(\infty,1)$, whose image is not contained in a hyperplane. In fact, by repeating the procedure we construct mappings

$$f\colon S^3\to Q(\infty,b) \tag{90}$$

for any finite b. Finally, by considering the form $r=\sin\big(\frac{\pi}{2}(|z_1|^2+|z_2|^2)\big)$ we obtain a mapping from S^3 to $Q(\infty,\infty)$.

As in the finite dimensional case, the situation changes when the source is a hyperquadric not equivalent to a sphere. If we now apply Lemma 6.2, we notice the following immediate consequence.

Corollary 7.1. *Let $a\geq 2$, $b\geq 1$. Let $U\subset Q(a,b)$ be a connected open set and $f\colon U\to Q(\infty,B)$, where $B\in\mathbb{N}_0\cup\{\infty\}$, be a real-analytic CR mapping such that $f(U)$ is not contained in any complex hyperplane of ℓ^2. Then $B=\infty$.*

Therefore, there is no mapping from $Q(a,b)$ for finite a and b into a $Q(\infty,B)$ for finite B. This rigidity result does not generalize to other hypersurfaces with indefinite nondegenerate Levi-form. In this case we can in fact construct mappings to $Q(\infty,B)$ for finite B.

For example, define r as

$$r(z,\bar z) = e^{|z_1+1|^2+|z_2|^2}-e-|z_3|^2. \tag{91}$$

By expanding the series we see that the signature pair of r is $(\infty,2)$. Furthermore, we look at the series up to the quadratic terms

$$r(z,\bar z) = 2e\operatorname{Re} z_1+|z_2|^2-|z_3|^2+\text{higher order terms.} \tag{92}$$

We see that the Levi form of $M=\{r=0\}$ at the origin is indefinite; that is, it has one positive and one negative eigenvalue. As the signature pair is $(\infty,2)$ we obtain a real-analytic

CR mapping

$$f \colon M \to Q(\infty, 1) \tag{93}$$

whose image is not contained in a hyperplane. Similarly we can easily obtain a submanifold with b negative eigenvalues and arbitrary number of positive eigenvalues in the Levi-form that admits a real-analytic CR mapping to $Q(\infty, b)$ whose image is not contained in a hyperplane.

The same construction can be used in the finite dimensional case. By truncating the series expansion of r, we obtain a polynomial r' and a manifold $M' = \{r' = 0\}$ with an indefinite Levi-form at 0, and such that r' has signature pair $(A, 2)$ for an arbitrarily high finite A. So we obtain a mapping from M' to $Q(A, 1)$ whose image is not contained in a hyperplane.

8. Construction of mappings

Let us give some explicit examples in nonhomogeneous coordinates before we give a general method for constructing mappings.

Example 8.1. Let $(z, w) \in \mathbb{C}^a \times \mathbb{C}^b$. We tensor with the identity (which we write as (z, w)) on the w_b variable. That is, we construct the mapping

$$(z, w) \mapsto \bigl(z, w_1, \ldots, w_{b-1}, w_b \otimes (z, w)\bigr). \tag{94}$$

We reorder the components to obtain a mapping taking $Q(a, b)$ to $Q(2a, 2b - 1)$. That is,

$$(z, w) \mapsto \bigl(z, w_b z_1, w_b z_2, \ldots, w_b z_a, w_1, w_2, \ldots, w_{b-1}, w_b w_1, w_b w_2, \ldots, w_b^2\bigr). \tag{95}$$

Therefore the super-rigidity result of Baouendi, Ebenfelt, and Huang [3] is sharp. In the same language as in the introduction we have that $A = 2a$ and $B = 2b - 1$. On the other hand, by Theorem 1.3 we are not able to make A arbitrarily large while keeping $B = 2b - 1$.

Example 8.2. We could repeat the procedure of the first example any number of times. We take a mapping $\varphi \colon Q(a, b) \to Q(A, B)$ and construct

$$\bigl(\varphi_1, \ldots, \varphi_{A+B-1}, \varphi_{A+B} \otimes (z, w)\bigr). \tag{96}$$

Again we reorder the components to obtain

$$\begin{split}\bigl(\varphi_1(z, w), \ldots, \varphi_A(z, w), z_1\varphi_{A+B}(z, w), \ldots, z_a\varphi_{A+B}(z, w), \\ \varphi_{A+1}(z, w), \ldots, \varphi_{A+B-1}(z, w), w_1\varphi_{A+B}(z, w), \ldots, w_b\varphi_{A+B}(z, w)\bigr)\end{split} \tag{97}$$

taking $Q(a, b)$ to $Q(A + a, B + b - 1)$.

For the constructions, it is easier to work in homogeneous coordinates. We define the homogeneous hyperquadric

$$HQ(a, b) := \Bigl\{ z \in \mathbb{C}^{a+b} : \sum_{j=1}^{a} |z_j|^2 - \sum_{j=a+1}^{a+b} |z_j|^2 = 0 \Bigr\}. \tag{98}$$

The set $HQ(a, b)$ defines a real hypersurface in the projective space $\mathbb{CP}^{a+b-1}$. By setting $z_{a+b} = 1$ we obtain $Q(a, b - 1)$ in $\mathbb{C}^{a+b-1}$. The homogeneous setting is more symmetric, although one has to be careful when converting back to the nonhomogeneous situation.

Suppose that $r(z, \bar{z})$ is a bihomogeneous polynomial and vanishes on $HQ(a, b)$. If r has signature pair (A, B) (that is A positive and B negative eigenvalues), then we find homogeneous polynomials $f \colon \mathbb{C}^{a+b} \to \mathbb{C}^A$ and $g \colon \mathbb{C}^{a+b} \to \mathbb{C}^B$ with linearly independent components such that

$$r(z, \bar{z}) = \|f(z)\|^2 - \|g(z)\|^2; \tag{99}$$

therefore the mapping $z \mapsto \big(f(z), g(z)\big)$ takes $HQ(a,b)$ to $HQ(A,B)$. By dividing by g_B and setting $z_{a+b} = 1$ we obtain a rational CR mapping $F\colon Q(a,b-1) \to Q(A,B-1)$. If the components of (f,g) are linearly independent then the image of F does not lie in a complex hyperplane.

Therefore when constructing examples we will consider homogenous polynomial mappings from $HQ(a,b)$ to $HQ(A,B)$ with linearly independent components.

9. Stability

We wish to show that as long as the number of positive and negative eigenvalues of the target hyperquadric grow in a comparable way, then nontrivial hyperquadric mappings always exist. We have the following stability result. We make the normalization $a \geq b$ for $HQ(a,b)$.

Theorem 9.1. *Fix integers $a \geq b \geq 2$. There exists an M ($M = a^2 + ab - 2a + 1$ suffices) such that when $A + B \geq M$, where $A, B \geq 2$ and*

$$\frac{B-b+1}{A} \geq \frac{b-1}{a} \qquad \text{and} \qquad \frac{A-b+1}{B} \geq \frac{b-1}{a}, \tag{100}$$

then there exists a homogeneous polynomial mapping $F\colon HQ(a,b) \to HQ(A,B)$ with linearly independent components. That is, there exists a rational CR mapping $\tilde{F}\colon Q(a,b-1) \to Q(A,B-1)$ whose image is not contained in a complex hyperplane.

We note that in [7] it is proved that when the source is a sphere (that is, $HQ(a,1)$) then such a rational mapping always exists for large $A+B$ provided only that $A,B \geq 1$. In the hyperquadric case, this sort of stability result holds in a sector, where the size of the sector depends on a and b. There may exist mappings outside of the sector given above; however, by Theorem 1.1 it is clear that any sector in which existence holds must make an acute angle.

The signature pairs of mappings that are constructed in the proof along with the inequalities are illustrated in Figure 1.

Proof. Suppose that we can find a homogeneous polynomial $p(x)$ in the variables $x_1, \ldots, x_{a+b}$ with A positive and B negative coefficients such that $p(x) = 0$ when $x_1 + \cdots + x_a - x_{a+1} - \cdots - x_{a+b} = 0$. We call such polynomials admissible for (A,B). We define

$$s := x_1 + \cdots + x_a - x_{a+1} - \cdots - x_{a+b}. \tag{101}$$

If p is admissible for (A,B) then we let $x_k = |z_k|^2$ in p. We obtain a Hermitian symmetric polynomial

$$r(z,\bar{z}) = p(x). \tag{102}$$

As distinct monomials are linearly independent, the rank of r is equal to the number of distinct monomials of p, that is $A+B$ and the signature pair is (A,B). As r vanishes on $s = 0$, we have that r induces a CR mapping of $HQ(a,b)$ to $HQ(A,B)$ such that the image is not contained in a complex hyperplane.

We therefore work with admissible polynomials p as above. We fix a and b and we work in the plane $\mathbb{N}_0^2$ and simply say $(A,B) \in E \subset \mathbb{N}_0^2$ whenever an admissible p exists for those (A,B) (a mapping to $HQ(A,B)$ exists).

We note that $(a,b) \in E$ and $(b,a) \in E$. If p is admissible for (A,B) of degree m then $\tilde{p} = x_1^{k-m} p + x_2^{k-1} s$ vanishes when s vanishes. If we pick k large enough then the monomials of $x_1^{k-m} p$ and $x_2^{k-1} s$ are distinct and $\tilde{p}$ is admissible for $(A+a, B+b)$. By using $-s$ instead



FIGURE 1. Diagram of mappings constructed in the proof of Theorem 9.1 for the source hyperquadric $HQ(4, 2)$. The lattice represents the positive quadrant of the (A, B) plane. A black dot represents that a mapping to $HQ(A, B)$ exists by applying the theorem. A black square represents that a mapping was also constructed in the proof. A circle represents a position in the lattice for which the theorem tells us nothing. The three lines represent the limits of the stability region from the theorem. The dashed squares represent one succession of the "squares" constructed in the proof.

of s we build an admissible polynomial for $(A + b, B + a)$. Therefore for any $n, k \in \mathbb{N}_0$ we have that

$$(a, b) + n(a, b) + k(b, a) \in E \qquad \text{and} \qquad (b, a) + n(a, b) + k(b, a) \in E. \tag{103}$$

Therefore we get a lattice of admissible points. We claim that for each point $(b, a) + n(a, b) + k(b, a)$ we also obtain that the "square"

$$\{(b, a) + n(a, b) + k(b, a) - (j, m) : 0 \leq j, m \leq n + k\} \tag{104}$$

is a subset of E. Once we have the claim, it is a simple matter of noticing that the squares must overlap for large enough $n + k$. The inequalities from the theorem follow. See Figure 1 for illustration of the proof.

To prove the claim, we need to show that if (A, B) is admissible, then the points

$$\begin{aligned} &(A, B) + (a, b), &&(A, B) + (a, b - 1), \\ &(A, B) + (a - 1, b), &&(A, B) + (a - 1, b - 1), \end{aligned} \tag{105}$$

and

$$\begin{aligned} &(A, B) + (b, a), &&(A, B) + (b, a - 1), \\ &(A, B) + (b - 1, a), &&(A, B) + (b - 1, a - 1). \end{aligned} \tag{106}$$

are also admissible.

The first point we have already seen belongs to E. Now take p that is admissible for (A, B). By dividing through by x_{a+b} we can suppose that p contains a monomial m that does not depend on x_{a+b}. Furthermore we can assume that m has a positive coefficient, since if no such monomial existed, setting $x_{a+b} = 0$ would obtain a contradiction.

Write $p = m + q$ and construct

$$\tilde{p} = m(s + x_{a+b}) + x_{a+b}q. \tag{107}$$

Where we note that $s + x_{a+b} = x_1 + \ldots - x_{a+b-1}$. The polynomial $\tilde{p}$ vanishes when s vanishes. As the coefficient of m is positive then $\tilde{p}$ is admissible for have $(A, B) + (a - 1, b - 1)$.

Similarly we can construct

$$\hat{p} = \frac{m(s + x_{a+b})}{2} + \frac{x_{a+b}m}{2} + x_{a+b}q. \tag{108}$$

The polynomial $\hat{p}$ vanishes when s vanishes. As the coefficient of m is positive then $\tilde{p}$ is admissible for have $(A, B) + (a, b - 1)$.

All the other points in the claim follow by variations on the above construction. □

References

[1] N. I. Akhiezer and I. M. Glazman, *Theory of linear operators in Hilbert space*, Dover Publications Inc., New York, 1993. MR1255973

[2] M. Salah Baouendi, Peter Ebenfelt, and Linda Preiss Rothschild, *Real submanifolds in complex space and their mappings*, Princeton Mathematical Series, vol. 47, Princeton University Press, Princeton, NJ, 1999. MR1668103

[3] M. S. Baouendi, Peter Ebenfelt, and Xiaojun Huang, *Holomorphic Mappings between Hyperquadrics with Small Signature Difference*, Amer. J. Math. (to appear). preprint arXiv:906.1235.

[4] M. S. Baouendi and Xiaojun Huang, *Super-rigidity for holomorphic mappings between hyperquadrics with positive signature*, J. Differential Geom. **69** (2005), no. 2, 379–398. MR2169869

[5] John P. D'Angelo, *Several complex variables and the geometry of real hypersurfaces*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1993. MR1224231

[6] John P. D'Angelo and Jiří Lebl, *Complexity results for CR mappings between spheres*, Internat. J. Math. **20** (2009), no. 2, 149–166. MR2493357

[7] ———, *Hermitian symmetric polynomials and CR complexity*, J. Geom. Anal. **21** (2011), no. 3, 599–619. MR2810845

[8] John P. D'Angelo, *Hermitian analogues of Hilbert's 17-th problem*, Adv. Math. **226** (2011), no. 5, 4607–4637. MR2770459

[9] John P. D'Angelo and Dror Varolin, *Positivity conditions for Hermitian symmetric functions*, Asian J. Math. **8** (2004), no. 2, 215–231. MR2129535

[10] Franc Forstnerič, *Extending proper holomorphic mappings of positive codimension*, Invent. Math. **95** (1989), no. 1, 31–61. MR969413

[11] ———, *Embedding strictly pseudoconvex domains into balls*, Trans. Amer. Math. Soc. **295** (1986), no. 1, 347–368, DOI 10.2307/2000160. MR831203

[12] Mark Green, *Restrictions of linear series to hyperplanes, and some results of Macaulay and Gotzmann*, Algebraic curves and projective geometry, 1989, pp. 76–86. MR1023391

[13] Xiaojun Huang, Shanyu Ji, and Dekang Xu, *A new gap phenomenon for proper holomorphic mappings from B^n into B^N*, Math. Res. Lett. **13** (2006), no. 4, 515–529. MR2250487

[14] Jiří Lebl and Daniel Lichtblau, *Uniqueness of certain polynomials constant on a line*, Linear algebra Appl. **433** (2010), 824–837. MR2654111

[15] László Lempert, *Imbedding Cauchy-Riemann manifolds into a sphere*, Internat. J. Math. **1** (1990), no. 1, 91–108, DOI 10.1142/S0129167X90000071. MR1044662

[16] Hans Lewy, *On the local character of the solutions of an atypical linear differential equation in three variables and a related theorem for regular functions of two complex variables*, Ann. of Math. (2) **64** (1956), 514–522. MR0081952

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA
E-mail address: `grundmer@umich.edu`

Department of Mathematics, University of Wisconsin, Madison, WI 53706, USA
E-mail address: `lebl@math.wisc.edu`

Department of Mathematics, Purdue University, West Lafayette, IN 47907, USA
E-mail address: `lvivas@math.purdue.edu`